

www.kadlecmed.org

healthy heart

A service of Kadlec Medical Center—Tri-Cities

HEALTH NEWS AND INFORMATION

cardiac care CLOSE TO HOME

IN JUST THE PAST TWO YEARS, more than 3,000 patients have received a diagnostic heart catheterization, more than 1,500 patients have had a balloon inflated or stents placed in their coronary arteries, and hundreds have had open-heart surgery at Kadlec Medical Center.

Because of Kadlec's superior cardiac team, people with heart disease can receive important and often lifesaving treatment close to home.

The program continues to grow with the arrival of a second cardiothoracic surgeon, Hannan Chaugle, MD, who joined the Kadlec Clinic cardiothoracic surgical team headed up by Juan Cordero Jr., MD.

Providing additional support are two cardiac physician assistants (PA-C), Jared Collet and April Derand. Derand is the only PA-C in

diagnosis through its chest pain clinic, angioplasty catheterization, open-heart surgery and cardiac rehabilitation. Kadlec is the only hospital in the region with this comprehensive scope of cardiac care.

Meeting your needs

"The Kadlec medical staff includes 12 cardiologists, three of whom are interventional cardiologists performing the balloon catheterizations and placement of stents in the catheterization lab," says Don Hanna, Cardiac Service Line program manager. "Our two cardiothoracic surgeons do the heart surgeries in Kadlec's new cardiovascular operating room."

"Our vision for cardiac services at Kadlec is focused on meeting the needs of our region. We know care close to home is the best for the patient and their family. If a patient from Hermiston can receive appropriate care in Hermiston, that's where they should stay. If they



Hannan Chaugle, MD, and Juan Cordero Jr., MD

need a higher level of care, we can provide it so they can get it as fast as possible yet remain as close to home as they can be," Hanna says.

Dr. Chaugle agrees. "It is our vision to create a center of excellence for cardiac care here at Kadlec. We want people in our region to look to us for full and comprehensive cardiac care when they need it," he says.

Kadlec has the most comprehensive scope of cardiac care in the region.

the region trained in endoscopic vein harvesting—the least invasive way to obtain veins for bypass surgery.

Kadlec performs the full spectrum of cardiac care, including fast

HEALTH facts

SPOUSES OF SMOKERS AT RISK

If you're a smoker married to a nonsmoker, do something that will help both of you: Quit. Nonsmokers who are married to smokers are at high risk of having a stroke. The good news: If you quit, you'll lower the risk that your spouse will have a stroke. And you'll help protect your health as well.

—American Journal of Preventive Medicine

WATCH OUT FOR ADDED SUGARS

One way to help control your weight is to cut back on beverages and foods with added sugars.

The calories in these products

can add up quickly. Examples of added sugars include sucrose, glucose, fructose, dextrose and high-fructose corn syrup. Read the ingredient list on labels, and choose items that don't have added sugars in their first four ingredients.

—American Heart Association



diuretics

WHAT YOU NEED TO KNOW



For more information about safe medicine use, visit Kadlec's health information library at www.kadlecmed.org.

SOMETIMES FLUID NEEDS A little help getting where it needs to go.

And in your body, a diuretic—or water pill—may be just the nudge your kidneys need to help excess fluid make an exit.

Diuretics work by helping your kidneys remove sodium and water from the bloodstream and convert them to urine. As that urine leaves your body (yes, via more frequent trips to the bathroom), it can:

- Help relieve the workload of your heart, since there's less fluid to pump throughout your body.
- Decrease the buildup of fluid in your lungs and other parts of your body, such as ankles and legs.

Besides frequent urination, diuretics can cause other side effects, including: ■ Dizziness. ■ Headache. ■ Thirstiness. ■ Muscle cramps. ■ Upset stomach.

To help get those pesky toilet trips out of the way during the day (instead of at night), many people take their diuretic in the morning.

A few cautions

Some types of diuretics remove potassium from the body, so you may need to take a supplement or eat more potassium-rich foods—such as bananas—or do both. Ask your doctor in order to be sure, though, because too much potassium can also be a problem.

Call your doctor if, while taking a diuretic, you develop: ■ A severe rash. ■ Trouble breathing or swallowing. ■ Gout, a condition that can cause pain and swelling in the joints (often the big toe). Your prescription may need to be adjusted.

Sources: American Heart Association; U.S. Food and Drug Administration



healthyheart

HEALTHY HEART is published as a community service for the friends and patrons of KADLEC MEDICAL CENTER, 888 Swift Blvd., Richland, WA 99352, 509-946-4611, website www.kadlecmed.org.

Rand Wortman
President and CEO, Kadlec Health System

Lane Savitch
President, Kadlec Medical Center

Jim Hall
Director, Community Relations

Information in HEALTHY HEART comes from a wide range of medical experts. If you have any concerns or questions about specific content that may affect your health, please contact your health care provider. Models may be used in photos and illustrations. Copyright © 2009 Coffey Communications, Inc. HTN23394h

legs

PROVIDE WARNING SIGNS OF PAD

YOU MAY THINK ONLY arteries that supply blood to the heart or brain can become blocked with plaque and cause problems.

But actually, arteries throughout the body can narrow and suffer reduced blood flow due to plaque. When that happens, the condition is known as peripheral arterial disease, or PAD. And the most common place for PAD to strike is in the arteries of the legs.

It's important to recognize warning signs of PAD and to get treatment for it, since it can put you at risk for heart attack, stroke and leg amputation.

Signs of PAD include: ■ Pain, cramping, fatigue or heaviness in the leg or hip muscles when you're active (such as walking or climbing stairs) but that goes away when you rest. ■ Sores on your feet or legs that don't heal well. ■ Pain, burning or aching in your legs or feet when you're resting. ■ Change in skin color on your legs. ■ Cooler spots on parts of your legs or feet.

Smoking is the No. 1 risk factor

PAD can put you at risk for heart attack and stroke.

for PAD, reports the American Medical Association. If you smoke, it is imperative that you stop; otherwise, treatment may not be effective.

Other risk factors include: age—PAD is more common after age 60; diabetes; high blood pressure; and high cholesterol.

Treating PAD

An essential part of treating PAD is adopting healthy habits. Don't smoke. Keep your blood pressure, cholesterol and blood sugar under control. Exercise regularly. Eat more whole grains, vegetables and fruit, and eat less salt, trans fat, saturated fat and cholesterol. And try to lose weight if you're overweight.

Your doctor may also prescribe medication to help prevent blood clots or to keep pain at bay when you're on the move. You may also need medication to keep diabetes, blood pressure or cholesterol in check.

More severely blocked arteries may require a procedure to open the artery or to bypass it.

Additional source: National Institutes of Health

Diagnosing PAD

If your doctor suspects that you have peripheral arterial disease, he or she will likely start with a few preliminary checks, such as:

- Asking about your medical history and any symptoms you have had.
- Examining your legs and feet for their pulse, color and temperature.
- Inspecting your lower limbs for any wounds that aren't healing well.

Other tests may be done as well, including:

Ankle-brachial index, or ABI.

For this test, the blood pressure in your ankle on one side of your body is compared to the pressure in your arm on the same side. Those two numbers should be roughly equal. If the pressure in your ankle is significantly lower, it's a sign that the arteries in your leg are probably narrowed.

Ultrasound. Sound waves are used to show the structure of your arteries as well as blood flow.

Angiogram. This test uses an injected dye and x-rays to check for narrowed or blocked arteries.

Source: Society for Vascular Surgery

KADLEC MEDICAL CENTER
888 Swift Blvd.
Richland, WA 99352

Nonprofit Org.
U.S. Postage
PAID
Tacoma, WA
Permit No. 1066



THE FACTS ABOUT fat

FAT ISN'T JUST FAT—THERE'S more to it than that.

Here's the skinny:

Saturated fat is the main dietary cause of high blood cholesterol. It's found mostly in animal products, such as high-fat cuts of meat, butter, cream, milk and cheeses. Certain plant products—palm oils, coconut oils and cocoa butter—also contain saturated fat.

Trans fats, or partially hydrogenated oils, are created during food processing when liquid oils are converted to solid fats. Trans fats are often found in commercially baked items, such as cookies, crackers and pies.

In the body, trans fats raise bad (LDL) cholesterol and lower good (HDL) cholesterol.

Unsaturated fats—when used in moderation and in place of saturated and trans fats—may actually help lower blood cholesterol.

There are two types: ■ Polyunsaturated fats are typically liquid at room temperature and when chilled. ■ Monounsaturated fats are typically liquid at room temperature but start to turn solid when chilled.

Unsaturated fats are found in fish such as salmon, mackerel and trout; nuts, such as walnuts and sunflower seeds; and vegetable oils, such as soybean, olive, canola, sesame, corn and safflower.

Most of the fat you eat should be unsaturated. But limiting your intake of all types of fat is best.

Sources: American Heart Association; U.S. Centers for Disease Control and Prevention



Artichoke salad with baby greens

Ingredients

- 1 cup of baby field greens
- 6 cherry tomatoes, halved
- ½ can (14 oz.) artichoke hearts, drained and cut into quarters*
- 1 tablespoon rice vinegar
- 2 teaspoons extra virgin olive oil
- 1 small clove garlic, finely minced
- Salt and freshly ground black pepper, to taste
- ½ tablespoon grated Parmesan cheese

*Remaining artichoke hearts may be refrigerated for up to five days or frozen for one to two months.

Instructions

1. In serving bowl, arrange greens, tomatoes and artichoke hearts.
2. In separate small bowl, whisk together vinegar, olive oil, garlic, salt and pepper. Toss into salad, and garnish with cheese.

Preparation time: 15 minutes

Nutrition information

Makes 2 servings.

Per serving: 91 calories, 5g total fat (<1g saturated fat), 9g carbohydrates, 4g protein, 2g dietary fiber, 318mg sodium

Source: Adapted from the American Institute for Cancer Research

Kadlec has nutrition counselors available by appointment at 509-942-2620.